

On the Dimension of The Virtually Cyclic Classifying Space of a Crystallographic Group

Frank Connolly, Benjamin Fehrman and Michael Hartglass

February 2, 2008

Abstract

In this paper we construct a model for the classifying space, $B_{\mathcal{VC}}\Gamma$, of a crystallographic group Γ of rank n relative to the family $\mathcal{VC}$ of virtually-cyclic subgroups of Γ . The model is used to show that there exists no other model for the virtually-cyclic classifying space of Γ with dimension less than $\text{vcd}(\Gamma)+1$, where $\text{vcd}(\Gamma)$ denotes the virtual cohomological dimension of Γ . In addition, the dimension of our construction realizes this limit.

1 Introduction. Statement of Results

Let Γ be a discrete group. Its classifying space, $B\Gamma = E\Gamma/\Gamma$, is a polyhedron for which the Γ -space $E\Gamma$ is a terminal object in the category of free Γ -CW-complexes and homotopy classes of Γ -maps. $B\Gamma$ is therefore unique up to homotopy type.

In recent years topologists and others (e.g. Brown [5], Farrell and Jones [9], Lueck [10] and Serre [11]), have found great use for the more general notion of a *classifying space of a discrete group Γ , relative to a family $\mathcal{F}$* , of subgroups of Γ . This is a space $B_{\mathcal{F}}\Gamma = E_{\mathcal{F}}\Gamma/\Gamma$, where $E_{\mathcal{F}}\Gamma$ is a Γ -CW complex which is a terminal object in the category of homotopy classes of Γ -maps between Γ -CW-complexes whose isotropy groups are in $\mathcal{F}$.

According to Eilenberg and Ganea [7], if Γ has cohomological dimension $\leq n$, then one can construct a model of $B\Gamma$ of dimension $\leq \max(n, 3)$. There are similar results showing $\text{hdim}(B_{\mathcal{FIN}}\Gamma) \leq \max(n, 3)$, in many cases (see

[6], [10]), but not all (see [4]), whenever $vcd(\Gamma) \leq n$. Here $\mathcal{FIN}$ denotes the family of finite subgroups of Γ , and we write

$$hdim(Y) = \min\{dim(X) \mid X \text{ is a CW-complex homotopy equivalent to } Y\}$$

A remarkable paper of F.T. Farrell and L. Jones [9] introduces the family

$$\mathcal{VC} = \{H \mid H \text{ is a virtually cyclic subgroup of } \Gamma\}$$

(A group is *virtually cyclic* if it contains a cyclic subgroup of finite index). They conjecture there that the K or L theory of a group Γ can be computed from the homology of $B_{\mathcal{VC}}\Gamma$, taken with stratified coefficients in the K or L theory of the virtually cyclic subgroups of Γ . The K-theoretic version of this conjecture has been proved, for many groups, by Bartels and Reich [2].

Farrell and Jones [9] also give two constructions of this space $B_{\mathcal{VC}}\Gamma$, one of which is a finite dimensional CW-complex when Γ is a discrete subgroup of a Lie Group.

The goal of this note is to provide a geometrically simple construction of $B_{\mathcal{VC}}\Gamma$, as an $n + 1$ -dimensional CW-complex, when Γ is a crystallographic group of rank n (see Section 3). We use this construction to prove the following result:

Theorem 1.1. Let Γ be a crystallographic group of rank $n \geq 2$. Then:

$$hdim(B_{\mathcal{VC}}\Gamma) = n + 1$$

Note that a crystallographic group Γ of rank 1 is virtually-cyclic. So in this case $hdim(B_{\mathcal{VC}}\Gamma) = 0$.

2 Basic Ideas.

A crystallographic group is a discrete co-compact subgroup of $\text{Iso}(\mathbb{R}^n)$, the group of isometries of $\mathbb{R}^n$. In this paper, the *translation subgroup* of a crystallographic group Γ will be denoted

$$A = \Gamma \cap \text{Trans}(\mathbb{R}^n),$$

where $\text{Trans}(\mathbb{R}^n)$ denotes the subgroup of $\text{Iso}(\mathbb{R}^n)$ consisting of translations. A is normal in Γ . The *point group* (or *holonomy group*) of Γ will be written

$$G = \Gamma/A.$$

The following theorem of Bieberbach [3] shows that G is a finite group.

Theorem 2.1. Let Γ be a crystallographic group of rank n . Then, A is a finitely generated, free abelian group of rank n with finite index in Γ .

A collection of subgroups $\mathcal{F}$ of a group Γ is called a *family* if $\mathcal{F}$ is closed under taking subgroups and under conjugation in Γ .

Definition 2.2. Let $\mathcal{F}$ be a family of subgroups of Γ . A Γ -space E is called $\mathcal{F}$ -universal if it satisfies the following conditions:

- i.) E^H is contractible. $\forall H \in \mathcal{F}$
- ii.) $E^H = \emptyset$. $\forall H \notin \mathcal{F}$
- iii.) E is a Γ -CW-complex

where E^H denotes the fixed set of a subgroup $H \subset \Gamma$ in E .

Then, we say E/Γ is a classifying-space for Γ relative to $\mathcal{F}$. By [10] this specifies E/Γ uniquely up to homotopy type.

We will mainly be interested in the family:

$$\mathcal{VC} = \{H \subset \Gamma \mid H \text{ a virtually-cyclic subgroup}\}.$$

Later, our model for a $\mathcal{VC}$ -universal space of a crystallographic group Γ will be denoted $E_{\mathcal{VC}}\Gamma$. The classifying space for Γ relative to $\mathcal{VC}$ will be written $B_{\mathcal{VC}}\Gamma = E_{\mathcal{VC}}\Gamma/\Gamma$.

Finally, let $\mathcal{C} = \{H \subset A \mid H \text{ is a maximal cyclic subgroup of } A\}$ be the set of maximal cyclic subgroups of A . For each subgroup $C \in \mathcal{C}$ we define:

$$\mathbb{R}^{n-1}(C) = \{l \subset \mathbb{R}^n \mid l \text{ a line, } C \cdot l = l\}.$$

This is the set of lines in $\mathbb{R}^n$ left invariant under the group action of C . The quotient map $\pi_C : \mathbb{R}^n \rightarrow \mathbb{R}^{n-1}(C)$ is the map:

$$\pi_C(x) = \text{the unique line } l \in \mathbb{R}^{n-1}(C) \text{ containing } x \quad \forall x \in \mathbb{R}^n.$$

The Hausdorff metric on the non-empty closed sets of $\mathbb{R}^n$ restricts to a metric on $\mathbb{R}^{n-1}(C)$. In addition, the quotient topology on $\mathbb{R}^{n-1}(C)$ coincides with the metric topology on $\mathbb{R}^{n-1}(C)$. Note that $\mathbb{R}^{n-1}(C)$ is isometric to $\mathbb{R}^{n-1}$ in such a way that π_C is a linear map.

3 Construction of $E_{\mathcal{V}\mathcal{C}}\Gamma$

Let Γ be a crystallographic group of rank n , with holonomy group G . Our model for $B_{\mathcal{V}\mathcal{C}}\Gamma$ will be the orbit space of a G -action on an infinite union of solid $n+1$ -tori sharing a common boundary. $E_{\mathcal{V}\mathcal{C}}\Gamma$ will be a similar union of mapping cylinders sharing a common source, one cylinder for each $C \in \mathcal{C}$.

Equip $\mathcal{C}$ with the discrete topology. Define an equivalence relation $\sim$ on $\mathbb{R}^n \times I \times \mathcal{C}$ as follows: $(x, t, C) \sim (x', t', C')$ if

- i.) $0 < t = t' < 1$, $C = C'$ and $x = x'$, or
- ii.) $t = t' = 1$, $C = C'$ and $\pi_C(x) = \pi_{C'}(x')$, or
- iii.) $t = t' = 0$ and $x = x'$.

We define:

$$E = (\mathbb{R}^n \times I \times \mathcal{C}) / \sim \quad (3.1)$$

with the quotient topology. The equivalence class of an element (x, t, C) is written $[x, t, C]$. For each $C \in \mathcal{C}$ we write

$$Cyl(\pi_C) = (\mathbb{R}^n \times I \times \{C\}) / \sim .$$

Note this subspace of E is just the mapping cylinder of $\pi_C : \mathbb{R}^n \rightarrow \mathbb{R}^{n-1}(C)$, and E is the union of these subspaces $Cyl(\pi_C)$.

The action of Γ on E is defined by:

$$\gamma \cdot [x, t, C] = [\gamma \cdot x, t, \gamma C \gamma^{-1}] \quad \forall \gamma \in \Gamma. \quad (3.2)$$

In the following sections we prove:

Proposition 3.1. The Γ -space E constructed in (3.1) and (3.2) is a $\mathcal{V}\mathcal{C}$ -universal Γ -space (see Definition 2.2).

4 E is a Γ -CW-Complex

In this section, we show that E has the structure of a Γ -CW-complex. Choose any Γ -CW-complex structure on $\mathbb{R}^n$, called X , whose cells, $e \in Cell(X)$ are all convex polytopes. Here, $Cell(X)$ is the collection of cells of X . It is enough to show that for each $C \in \mathcal{C}$, this CW-structure extends

equivariantly to a CW-structure on $Cyl(\pi_C)$. This would be obvious if $\pi_C: X \longrightarrow \mathbb{R}^{n-1}(C)$ were cellular with respect to a CW-complex structure Y on $\mathbb{R}^{n-1}(C)$, for we would then use the resulting CW-complex structure on $Cyl(\pi_C)$. It is nearly as obvious if there is a subdivision X' of X (that is, each cell of X' will be a subset of a cell of X) relative to which $\pi_C: X' \longrightarrow Y$ is cellular, since $X \times I$ can be subdivided by replacing only the cells of $X \times \{1\}$ with the cells of $X' \times \{1\}$ and leaving the other cells unchanged. This means that it suffices to show that $\mathbb{R}^{n-1}(C)$ has the structure of an $N_\Gamma(C)$ -CW-complex so that π_C is cellular relative to the subdivision X' of X .

First note $\{\pi_C(e) \mid e \in Cell(X)\}$ is a locally finite collection in $\mathbb{R}^{n-1}(C)$, because $Cell(X)$ is a locally finite collection in $\mathbb{R}^n$. We note that a polytope generated by n points is the convex hull of those n points, and we choose X so that $Cell(X)$ is generated by convex polytopes. For each $y \in \mathbb{R}^{n-1}(C)$ define:

$$f_y = \bigcap \{\pi_C(e) \mid y \text{ is contained in the interior of the convex polytope } \pi_C(e)\}.$$

This intersection is finite, so we have

$$Int(f_y) = \bigcap \{Int(\pi_C(e)) \mid y \in Int(\pi_C(e))\}.$$

Note $y \in Int(f_y)$ so each point of $\mathbb{R}^{n-1}(C)$ is in the interior of exactly one of the convex polytopes f_y . Also, $\{f_y \mid y \in \mathbb{R}^{n-1}(C)\}$ is a locally finite collection. To see that these form an $N_\Gamma(C)$ -CW-complex we show that if $dim(f_y) = d$, then $\dot{f}_y$ is a union of cells of dimension $< d$ (here, $\dot{f}_y$ denotes the boundary of f_y). Let $y' \in \dot{f}_y$. Then $y' \in \pi_C(e)$ for some e such that $y \in Int(\pi_C(e))$. Note that $\pi_C(e)^\circ$ is a subcomplex of $\pi_C(\dot{e})$. But $dim(\pi_C(e)) = d$, so $\pi_C(e)^\circ$ is a union of cells $\pi_C(e')$ of dimension $\leq d - 1$. Hence, there is a cell $\pi_C(e')$ with $y' \in Int(\pi_C(e'))$ such that $\pi_C(e') \subset \pi_C(e)^\circ$. Therefore, $\dot{f}_y$ is a union of cells of dimension $< d$. This shows that $Cell(Y) := \{f_y \mid y \in \mathbb{R}^{n-1}(C)\}$ is the collection of cells of an $N_\Gamma(C)$ -CW-complex structure on $\mathbb{R}^{n-1}(C)$ denoted Y . The cells of X' are now easily defined. They are the nonempty sets of the form $e \cap \pi_C^{-1}(f)$ with $e \in Cell(X)$ and $f \in Cell(Y)$. Note that $e \cap \pi_C^{-1}(f)$ is a convex polytope contained in the cell e . Since π_C takes each cell into a cell linearly, it follows that $\pi_C: X' \longrightarrow Y$ is cellular. The above observations show that E admits the structure of a Γ -CW-complex.

5 Contractibility of Fixed Sets

In this section we prove:

Lemma 5.1. Let E be the Γ -space constructed in (3.1). For each subgroup $H \subset \Gamma$, E^H is contractible if $H \in \mathcal{VC}$ and E^H is empty if $H \notin \mathcal{VC}$.

Proof. Let H be a subgroup of Γ . Then,

$$(\mathbb{R}^n)^H \text{ is } \begin{cases} \text{contractible} & \text{if } |H| < \infty \\ \emptyset & \text{if } |H| = \infty. \end{cases} \quad (5.1)$$

The Γ -action of (3.2) defines an action of $N_\Gamma(C)/C$ on $\mathbb{R}^{n-1}(C)$. This gives a map $j_C : N_\Gamma(C)/C \rightarrow \text{Iso}(\mathbb{R}^{n-1}(C))$. $\text{Ker}(j_C)$ has at most two elements. Recall $\mathbb{R}^{n-1}(C)$ is isometric to $\mathbb{R}^{n-1}$. The image of j_C is a crystallographic group.

Assume H is a subgroup of $N_\Gamma(C)$ for some $C \in \mathcal{C}$. By the above,

$$(\mathbb{R}^{n-1}(C))^H \text{ is } \begin{cases} \text{contractible} & \text{if } |HC/C| < \infty \\ \emptyset & \text{if } |HC/C| = \infty. \end{cases} \quad (5.2)$$

Claim 5.2. Let $H \subset \Gamma$ satisfy $|H| = \infty$. Then, there exists at most one $C \in \mathcal{C}$ satisfying $E^H \cap \text{Cyl}(\pi_C) \neq \emptyset$.

Proof. Suppose there are two subgroups $C, C' \in \mathcal{C}$ satisfying $E^H \cap \text{Cyl}(\pi_C) \neq \emptyset$ and $E^H \cap \text{Cyl}(\pi_{C'}) \neq \emptyset$. By (3.2) this implies $H \subset N_\Gamma(C)$ and $H \subset N_\Gamma(C')$. Then, by (5.1) and (5.2) we know $|HC/C| < \infty$ and $|HC'/C'| < \infty$. Therefore, the relation $HC/C \approx H/(H \cap C)$ shows that $H \cap C$ and $H \cap C'$ have finite index in H . So $H \cap C \cap C'$ has finite index in H . But $H \cap C \cap C' = \{e\}$ and $|H| = \infty$, a contradiction. $\square$

Claim 5.3. Let $H \subset \Gamma$ be an infinite subgroup. Then E^H is contractible if $H \in \mathcal{VC}$ and E^H is empty if $H \notin \mathcal{VC}$.

Proof. If $H \in \mathcal{VC}$ there exists precisely one $C \in \mathcal{C}$ satisfying $E^H \cap \text{Cyl}(\pi_C) \neq \emptyset$ (namely, the unique element of $\mathcal{C}$ containing $H \cap A$). For this C , (5.1) implies $E^H = (\mathbb{R}^{n-1}(C))^H \times \{1\} \times \{C\}$. This space is contractible by (5.2). Conversely, suppose there exists a unique $C \in \mathcal{C}$ satisfying $E^H \cap \text{Cyl}(\pi_C) \neq \emptyset$. Then, by (5.1) and (5.2) we know $H \in \mathcal{VC}$ because $|H/(H \cap C)| < \infty$. Therefore, E^H is contractible. Finally, if there exists no $C \in \mathcal{C}$ satisfying $E^H \cap \text{Cyl}(\pi_C) \neq \emptyset$ then $E^H = \emptyset$. $\square$

Thus, for a subgroup $H \subset \Gamma$ the set E^H is the following:

$$E^H = \begin{cases} (\mathbb{R}^n)^H \cup (\cup \{Cyl(\pi_C^H) \mid H \subset N_\Gamma(C)\}) & |H| < \infty \\ (\mathbb{R}^{n-1}(C))^H & |H| = \infty, H \in \mathcal{VC}, H \subset N_\Gamma(C) \\ \emptyset & |H| = \infty, H \notin \mathcal{VC}. \end{cases} \quad (5.3)$$

If $H \in \mathcal{VC}$ and H is infinite then E^H is contractible by Claim 5.3. If H is finite, then π_C^H is a homotopy equivalence for each $C \in \mathcal{C}$ for which $H \subset N_\Gamma(C)$ (here, π_C^H is the restriction of π_C to $(\mathbb{R}^n)^H$). Therefore the subcomplex $(\mathbb{R}^n)^H$ is a strong deformation retract of the CW-complex $Cyl(\pi_C^H)$, and these deformations coalesce to make $(\mathbb{R}^n)^H$ a strong deformation retract of E^H . This and (5.1) show that E^H is contractible. $\square$

Proof. (of Proposition 3.1): This is clear from Section 4 and Lemma 5.1. $\square$

Henceforth we denote the $\mathcal{VC}$ -universal Γ -space E by:

$$E = E_{\mathcal{VC}}\Gamma$$

and

$$E/\Gamma = B_{\mathcal{VC}}\Gamma.$$

6 Geometric Properties of $B_{\mathcal{VC}}\Gamma$

Let Γ , A and G be as in Section 2. In this section we will show that $B_{\mathcal{VC}}\Gamma$ is the orbit space of a G -action on an infinite union of solid $n+1$ -tori sharing a common boundary.

First we exhibit a homeomorphism

$$f : Cyl(\pi_C)/A \rightarrow D^2 \times T^{n-1}. \quad (6.1)$$

This will show $Cyl(\pi_C)/A$ is homeomorphic to the solid $n+1$ -torus $D^2 \times T^{n-1}$ for each $C \in \mathcal{C}$. Choose a basis $\{a_1, \dots, a_n\}$ for A so that a_1 generates C . Set $v_i = a_i(0)$. Then $\{v_1, \dots, v_n\}$ is a basis for $\mathbb{R}^n$. If $p = x_1 v_1 + \dots + x_n v_n$, we will write $p = \langle x_1, \dots, x_n \rangle$.

Let $f : Cyl(\pi_C)/A \rightarrow D^2 \times T^{n-1}$ be the map:

$$f([p, t, C]) = (1-t)e^{2\pi i x_1} \times e^{2\pi i x_2} \times \dots \times e^{2\pi i x_n},$$

where $p = \langle x_1, \dots, x_n \rangle$ and $[[p, t, C]]$ denotes the orbit, in $Cyl(\pi_C)/A$, of a point $[p, t, C] \in Cyl(\pi_C)$. The map f is continuous and bijective. It is a homeomorphism as $Cyl(\pi_C)/A$ is compact.

Because $E_{\mathcal{V}\mathcal{C}}\Gamma/A = \cup\{Cyl(\pi_C)/A \mid C \in \mathcal{C}\}$ the space $E_{\mathcal{V}\mathcal{C}}\Gamma/A$ is homeomorphic to an infinite union of solid $n+1$ -tori sharing the common boundary $\mathbb{R}^n/A$. Note that $\mathbb{R}^n/A$ is homeomorphic to T^n .

The relation $E_{\mathcal{V}\mathcal{C}}\Gamma/\Gamma \simeq (E_{\mathcal{V}\mathcal{C}}\Gamma/A)/(\Gamma/A)$ proves

$$B_{\mathcal{V}\mathcal{C}}\Gamma = (E_{\mathcal{V}\mathcal{C}}\Gamma/A)/G.$$

$B_{\mathcal{V}\mathcal{C}}\Gamma$ is therefore the orbit space of a G -action on a union of solid $n+1$ -tori sharing a common boundary, one torus for each $C \in \mathcal{C}$.

7 Computation of $hdim(B_{\mathcal{V}\mathcal{C}}\Gamma)$

In this section we prove Theorem 1.1.

Proof. Let $i_C: Cyl(\pi_C)/A \rightarrow E_{\mathcal{V}\mathcal{C}}\Gamma/A$ be inclusion, $\tau: E_{\mathcal{V}\mathcal{C}}\Gamma/A \rightarrow B_{\mathcal{V}\mathcal{C}}\Gamma$ be the obvious projection and $\iota_C = \tau \circ i_C: Cyl(\pi_C)/A \rightarrow B_{\mathcal{V}\mathcal{C}}\Gamma$. We will make a computation in the $n+1$ st homology group $H_{n+1}(B_{\mathcal{V}\mathcal{C}}\Gamma)$ using the cellular chain complex, $C_*(B_{\mathcal{V}\mathcal{C}}\Gamma)$. By (6.1), for each $C \in \mathcal{C}$ there exists an $n+1$ chain, $\chi_C \in C_{n+1}(Cyl(\pi_C)/A)$ such that $\partial_{n+1}(\chi_C) = [T_n]$ (where ∂_{n+1} is the boundary map and $[T_n]$ is the fundamental cycle of $\mathbb{R}^n/A$). Choose $C, C' \in \mathcal{C}$ such that C and C' are not conjugate in Γ (here is where we use that $rank(\Gamma) \geq 2$). Denote $\iota_{C\#}$ and $\iota_{C'\#}$ as the chain maps induced from ι_C and $\iota_{C'}$ respectively. Then, in $C_{n+1}(B_{\mathcal{V}\mathcal{C}}\Gamma)$:

$$z = \iota_{C\#}(\chi_C) - \iota_{C'\#}(\chi_{C'}) \neq 0. \text{ and } \partial_{n+1}(z) = 0.$$

Indeed,

$$\begin{aligned} \partial_{n+1}(z) &= \partial_{n+1}\iota_{C\#}(\chi_C) - \partial_{n+1}\iota_{C'\#}(\chi_{C'}) \\ &= \iota_{C\#}\partial_{n+1}(\chi_C) - \iota_{C'\#}\partial_{n+1}(\chi_{C'}) = \iota_{C\#}[T_n] - \iota_{C'\#}[T_n] = 0 \end{aligned}$$

as the restrictions of ι_C and $\iota_{C'}$ to $\mathbb{R}^n/A$ are the same. Since $C_{n+2}(B_{\mathcal{V}\mathcal{C}}\Gamma)$ is trivial, we conclude that $H_{n+1}(B_{\mathcal{V}\mathcal{C}}\Gamma)$ is nontrivial, implying that $hdim(B_{\mathcal{V}\mathcal{C}}\Gamma) \geq n+1$. But $B_{\mathcal{V}\mathcal{C}}\Gamma$ has dimension $n+1$. Therefore $hdim(B_{\mathcal{V}\mathcal{C}}\Gamma) = n+1$. $\square$

8 Conclusion and an Open Question

In this paper, we were able to construct a model for both $E_{\mathcal{VC}}\Gamma$ and $B_{\mathcal{VC}}\Gamma$ for a rank- n crystallographic group Γ . Despite the fairly straightforward constructions of our models, they are not locally finite CW-complexes. Therefore they can not be imbedded in any Euclidian space as they are not metrizable.

One question that is still unsolved is the following:

Conjecture 8.1. Suppose Γ is a discrete subgroup of a Lie Group and Γ is not virtually-cyclic. Then, $hdim(B_{\mathcal{VC}}\Gamma) = vcd(\Gamma) + 1$.

We have shown this is so if Γ is a crystallographic group, of rank $n \geq 2$.

References

- [1] Armstrong, M. A. *Basic Topology*. New York: Springer, 1983.
- [2] Bartels, Arthur and Holger Reich. "On the Farrell-Jones conjecture for higher algebraic K -theory." *J. Amer. Math Soc.* 18 (2005), no. 3, 501-545.
- [3] Bieberbach, L. "Über die Bewegungsgruppen der Euklidischen Raume II." *Math. Ann.* 72 (1912), 400-412.
- [4] Brady, Noel and Ian J. Leary and Brita E.A. Nucinkis. "On algebraic and geometric dimensions for groups with torsion." *J. London Math Soc.* (2) 64 (2001), no. 2, 489-500.
- [5] Brown, K.S. "Groups of Virtually Finite Dimension." *Homological Group Theory (C.T.C. Wall ed.)*. London Math Society Lecture Notes 36, Cambridge Univ. Press, Cambridge, 1979, 27-70.
- [6] Connolly, Frank and Tadeusz Kozniowski. "Finiteness Properties of Classifying Spaces of Proper Γ Actions." *J. Pure Appl. Algebra* 41 (1986), no. 1, 17-36.
- [7] Eilenberg, Samuel and Tudor Ganea. "On the Lusternik-Schnirelmann category of abstract groups." *Annals of Mathematics* (2) 65 (1957), 517-518.
- [8] Farkas, Daniel R. "Crystallographic Groups and Their Mathematics." *Rocky Mountain Journal of Mathematics* 11 (1981), no. 4, 511-551.
- [9] Farrell, F.T. and L.E. Jones. "Isomorphism Conjectures in Algebraic K -Theory." *Journal of the American Mathematical Society* 6 (1993), no.2, 249-297.
- [10] Lueck, Wolfgang. "The type of the classifying space for a family of subgroups." *Journal of Pure and Applied Algebra* 149 (2000), 177-203.
- [11] Serre, J.P. "Cohomologie des Groups Discrets." *Ann. of Math Studies* 70 (1971), 77-169.